Review

CellPress

To look or not to look: dissociating presaccadic and covert spatial attention

Hsin-Hung Li ,^{1,2,*} Nina M. Hanning ,^{1,2} and Marisa Carrasco ^{1,*}

Attention is a central neural process that enables selective and efficient processing of visual information. Individuals can attend to specific visual information either overtly, by making an eye movement to an object of interest, or covertly, without moving their eyes. We review behavioral, neuropsychological, neurophysiological, and computational evidence of presaccadic attentional modulations that occur while preparing saccadic eye movements, and highlight their differences from those of covert spatial endogenous (voluntary) and exogenous (involuntary) attention. We discuss recent studies and experimental procedures on how these different types of attention impact visual performance, alter appearance, differentially modulate the featural representation of basic visual dimensions (orientation and spatial frequency), engage different neural computations, and recruit partially distinct neural substrates. We conclude that presaccadic attention and covert attention are dissociable.

Presaccadic and covert attention are dissociable

Much of human visual experience results from moving our gaze to actively explore the visual world and gather information. By scanning the scene with **saccadic eye movements** (see Glossary), we bring relevant objects into our **fovea**, where visual information is processed with high precision. The link between eye movements and visual perception is so tight that perception is already facilitated before our gaze has reached a location of interest: immediately before an eye movement, while we prepare to saccade, presaccadic attention is deployed to the future gaze location, improving performance and altering stimulus appearance at that location.

Attention can also be deployed covertly toward specific locations, without moving the eyes. There are two types of covert attention. Exogenous attention is involuntary, stimulus-driven (bottom-up), and is transiently deployed within ~100 ms. By contrast, endogenous attention is voluntary, goal-driven (top-down), and deployed in a slower but sustained manner, typically reaching its full effect by ~300 ms [1,2]. Each type of covert attention improves performance and alters stimulus appearance at the attended location.

At a broader scale, the behavioral consequences and the neural correlates of presaccadic and covert attention appear to be similar. This resemblance led to the notion that covert attention may be a cognitive process subsidiary to oculomotor programming. Over a decade ago some studies questioned such a proposal. Given advances in experimental design and characterization of the attentional effects on performance and visual representation, recent studies have revealed novel dissociations between presaccadic and covert attention.

Highlights

To effectively process visual information, individuals attend to relevant visual inputs by selectively processing information at some locations at the expense of information elsewhere. One can attend overtly, by making saccadic eye movements, or covertly, by prioritizing information without moving the eyes. Both processes improve visual performance, but have different temporal dynamics.

Behavioral evidence shows that covert spatial attention is not the consequence of oculomotor planning.

Presaccadic and covert exogenous (involuntary) and endogenous (voluntary) attention differentially alter the featural representation of basic dimensions – orientation and spatial frequency.

Different neural computations mediate the effects of presaccadic and covert attention on contrast sensitivity.

Partially overlapping brain areas underlie presaccadic and covert attention, but different neuronal populations within each area subserve each type of attention.

¹Department of Psychology and Center for Neural Science, New York University, New York, NY, USA ²These authors contributed equally

*Correspondence: hsin.hung.li@nyu.edu (H.-H. Li) and marisa.carrasco@nyu.edu (M. Carrasco).



We bring together behavioral, neuropsychological, neurophysiological, and computational evidence distinguishing the effects of presaccadic attention[†] on visual performance and on featural representation from those of covert endogenous and exogenous attention.

Effects of presaccadic attention on visual performance

Linking saccadic eye movements and visual attention

Over the past three decades, different groups have found that perceptual judgments immediately before eye movement onset are more accurate for stimuli presented at the saccade target (the upcoming eye fixation) than elsewhere [5–7]. Interest in the perceptual consequences of presaccadic attention has continuously increased, and numerous studies have explored its spatial and temporal properties using variations of a typical dual-task protocol (Table 1).

Spatial and temporal coupling of oculomotor planning and attentional orienting

By testing performance at different locations relative to the saccade target and/or at different times relative to eye movement onset, behavioral studies – with varying spatial and temporal resolution – have provided consistent evidence of a tight spatial and temporal link between oculomotor programming and attentional orienting (Table 1). Spatially, perceptual benefits are typically bound to the saccade target and do not spread to neighboring stimuli [5,7,8], and the specific distribution of presaccadic attention is shaped by the spatial configuration of the scene [9–11]. Temporally, presaccadic attention builds up during saccade preparation, gradually reaching its maximum within ~75 ms before saccade onset, which is within ~200 ms after the cue indicating the saccade goal [8,12–19]. Whereas the close linkage between oculomotor and attentional orienting is undisputed, the nature of this coupling, and whether it is mandatory, has long been debated ([20] for recent review).

Is the coupling mandatory?

Early presaccadic attention studies claimed that eye movements and attention are mandatorily coupled – in other words, shortly before saccade onset one cannot attend elsewhere than to the saccade target [5–7] – a notion supported by the observation of worsened visual performance at non-saccade target locations. Although visual performance can be maintained to some degree at locations other than the saccade target during the early stages of movement planning, performance at non-target locations necessarily decreases just before saccade onset [13,21]. In line with this presaccadic cost at non-saccade targets [22,23], studies have demonstrated that attention is deployed to the saccade target even when detrimental for the task [7,24], suggesting a strong functional coupling between eye movements and attention.

Associations

The premotor theory of attention [25] even equated oculomotor and attentional orienting. Based on the observation of relatively prolonged reaction times for detecting targets appearing in the other visual hemifield than a spatial pre-cue, the authors postulated that a saccadic motor plan is required for the (re)orienting of visual spatial attention (because reorienting the eyes across the midline would also take longer as it requires the activation of different sets of muscles). That is, covert shifts of attention are achieved by programming an eye movement, without executing it. Accordingly, spatial attention and oculomotor processes are considered to rely on the same neural substrates; any shift of spatial attention – overt or covert – would be elicited by preceding motor activity in the oculomotor system.

⁺ We focus on perceptual modulations preceding saccadic eye movements (perceptual effects during the eye movement are reviewed in [3] and modulations specific to the remapped location in [4]).

Glossary

Adaptation: reduced sensory responses to prolonged or repeated presentations of the same stimulus. In psychophysical experiments, visual performance for a test stimulus following an adaptor stimulus is lower when the stimuli share similar features (e.g., orientation).

Antisaccade task: participants are required to suppress a reflexive saccade to a salient visual stimulus and instead perform a voluntary saccade in the opposite direction.

Fovea: a depression in the inner retinal surface, ~1.5 mm in width, whose photoreceptor layer entirely consists of cones and which is specialized for highest visual acuity.

Frontal eye field (FEF): in macaque monkeys, the FEF occupies the rostral bank of the arcuate sulcus; its putative human homolog is located near the precentral sulcus and the dorsal-most portion of the superior frontal sulcus. It plays an important role in guiding attention and saccades.

Global saccade: a saccade that involuntarily lands between two potential saccade targets presented at nearby locations.

Moebius syndrome: a rare congenital neurological condition, characterized by oculomotor disorders including gaze paralysis, that is caused by hypoplasia of the lower brainstem containing cranial nerves involved in the execution of eye movements.

Point of subjective equality (PSE): in comparative judgment experiments, the level (e.g., contrast) of the test stimulus varies while the standard stimulus remains constant. PSE is the test stimulus level at which both stimuli are chosen with equal probability and are judged as appearing equal.

Population receptive field (pRF): the spatial region of the visual field, usually modeled as a 2D Gaussian, that drives the response of a voxel in fMRI recording.

Reverse correlation: a technique for identifying the image features that drive sensory responses by presenting random (or pseudo-random) noise to the participants and correlating the image features with behavioral or neural responses.

Saccadic adaptation: an experimental procedure to dissociate saccade target and saccade landing location. By systematically displacing the target during the eye movement, the saccadic



Table 1. Studies and experimental procedures investigating the spatial and temporal coupling between saccadic eye movements and visual attention^a

Task/protocol	Standard	Temporal focus	Spatial focus
Discrimination task	Kowler <i>et al.</i> 1995 [5] Hoffman and Subramaniam 1995 [6] Deubel and Schneider 1996 [7] Montagnini and Castet 2007 [21] Zhao <i>et al.</i> 2012 [22] ★ Khan <i>et al.</i> 2015 [23] ★Li <i>et al.</i> 2019 [54] Kreyenmeier <i>et al.</i> 2020 [24] ▲ Parker <i>et al.</i> 2020 [148] ★ Li <i>et al.</i> 2021 [99]	Castet <i>et al.</i> 2006 [12] Deubel 2008 [13] Filali-Sacouk <i>et al.</i> 2010 [14] *Rolfs and Carrasco 2012 [15] White <i>et al.</i> 2013 [17] Harrison <i>et al.</i> 2013 [16] *Li <i>et al.</i> 2016 [18] Jonikaitis <i>et al.</i> 2017 [19] Hanning <i>et al.</i> 2019 [8]	▲ Ditterich <i>et al.</i> 2000 [38] Doré-Mazars and Collins 2005 [39] Ghahghaei and Verghese 2017 [9] Puntirol <i>et al.</i> 2018 [10] Szinte <i>et al.</i> 2019 [11]
Eye-abduction protocol	Smith <i>et al.</i> 2010 [28] Smith <i>et al.</i> 2012 [29] ▲ Hanning and Deubel 2020[35]		▲ Hanning <i>et al.</i> 2019 [34]
Global saccade task	▲ Van der Stigchel and de Vries 2015 [41]	▲ Wollenberg <i>et al.</i> 2020 [43]	▲ Wollenberg <i>et al.</i> 2018 [42]
Antisaccade task	▲ Klapetek <i>et al.</i> 2016 [44] Mikula <i>et al.</i> 2018 [45]		

^aKey: \blacktriangle , dissociations between attentional and oculomotor orienting; \bigstar , dissociation between overt and covert attention.

Additional studies supporting a dependence of visual attention on the oculomotor system rely on restricted ability to execute eye movements, either due to a pathology of the eye movement system (e.g., [26,27]) or experimentally induced by a maximally rotated eye position that prevents healthy participants from moving their eyes to particular peripheral locations (Figure 1B) [28,29]. According to both lines of research, although endogenous visual spatial attention can be properly deployed toward locations beyond the reach of the eyes, exogenous attention cannot [30].

However, the vast majority of studies claiming evidence in favor of the premotor theory (e.g., [27,29]) (as well as some studies arguing against it; e.g., [31–33]) have inferred attentional allocation from manual reaction times, following the rationale that stimuli are processed faster when they are presented within rather than outside the focus of attention. Nonetheless, reaction times reflect the combined effect of detection-, decision-, criterion- and response-dependent processes, which can only be differentiated with special methods (e.g., speed–accuracy trade-off) and models (e.g., drift-diffusion model) that are not used in these studies. Studies measuring discrimination performance can isolate these components and better assess attention effects because they can rule out speed–accuracy trade-offs and/or rely on signal detection theory (SDT) which indexes sensitivity and criterion separately [1,2].

Dissociations

Discrimination protocols have revealed that exogenous attention can be deployed at locations that the eyes cannot reach. A salient visual cue equally attracts attention regardless of whether it was presented within or beyond the reach of the eyes, causing perceptual benefits at its location [34] and perceptual costs elsewhere [35]. These studies, as well as reports that pathological oculomotor restrictions (e.g., **Moebius syndrome**) are not necessarily associated with corresponding attentional deficits [36], have revealed that covert attention can shift over the entire visual field range, independent of limitations imposed by the eye movement system [37].

amplitude and/or direction gradually adapts to the intra-saccadic shift. **Saccadic eye movements:** fast, ballistic movements of the eyes that shift gaze quickly toward a visual target.

Superior colliculus (SC): a structure in the midbrain, the SC is part of the brain circuit for the transformation of sensory input into movement output. It receives direct projections from retinal ganglion cells and conveys information to V1 and to extrastriate visual areas. It plays a major role in the control of eye movements and is involved in the control of covert attention.

Trans-saccadic integration: the integration of the presaccadic (e.g., a peripheral saccade target) and postsaccadic (e.g., the foveal image after saccade landing) information to facilitate visual stability across saccadic eye movements.

Visual area V1: located in the calcarine sulcus in the medial occipital lobe, V1 is the first visual processing area in the cortex. It codes for stimulus orientation, spatial frequency, and contrast. Visual area V4: a cortical area in the visual ventral stream that receives strong feedforward input from V2 and V1, and

sends connections to the inferior temporal cortex.

Visual masking: a phenomenon in which the visibility of a visual target is reduced by the presence of other visual stimuli (noise or mask) which overlap with or have a close proximity to the target in space and time.





Figure 1. Prominent experimental procedures to investigate the effects of presaccadic attention on visual performance. (Top row) Depiction of the respective task (saccade targets are indicated by green arrows, saccade endpoints are indicated by broken green lines). (Bottom row) Exemplary results. (A) Discrimination task. Participants prepare a saccade to an instructed saccade target (here the central arrow cue indicates the orange item on the right). A test stimulus ('E') is presented briefly before eye movement onset together with several distractors ('2's and '5's) at a location coinciding with the saccade target or not. Participants are asked to discriminate the test stimulus ('E' versus 'mirror-E'). While the eyes are still fixating the center, discrimination accuracy is selectively enhanced at the respective saccade target – indicating a presaccadic shift of attention. Adapted from [7]. (B) Eye abduction protocol. Participants view the display with one eye patched and their head rotated such that a portion of the display is still visible but outside the range reachable by the eyes (indicated by the green area). Saccades aimed at a cued target outside this range will fall short. Presaccadic discrimination accuracy (assessed with an oriented 1/f noise patch) is nonetheless highest at the unreachable saccade target (and not enhanced at the actual saccade landing) - demonstrating that attention is not limited to the eye movement range. Adapted from [34]. (C) Global saccade protocol. Participants are instructed to saccade to one of two nearby items at free choice (either of the orange frames), in which case the eyes will frequently land between the two salient targets. Presaccadic discrimination accuracy (assessed with an oriented Gabor patch) is not enhanced at the endpoint of such global saccades, but is instead equally enhanced at the two targets - showing that presaccadic attention is not necessarily coupled to the saccade endpoint. Adapted, with permission, from [42]. (D) Antisaccade task. Participants are instructed to saccade to the location in the opposite direction of a peripheral cue. Presaccadic discrimination accuracy (assessed with a tilted Gabor patch) before antisaccades is equally enhanced at the cued location and the antisaccade goal, indicating attentional selection at both locations. Adapted from [44]. (E) Microstimulation protocol. Monkeys must detect a luminance change of a peripheral target. Subthreshold frontal eye field stimulation increases visual sensitivity at the movement field (to which saccades would be evoked by stimulation currents above threshold; marked by the green circle), suggesting that saccade-related mechanisms provide a source of spatial attention. Adapted from [107]. Abbreviations: FEF, frontal eye field; SL, saccade landing position; ST, saccade target.

This decoupling of attentional and oculomotor orienting is in line with studies evaluating presaccadic attention as a function of the saccade endpoint. Even when saccades undershoot their target, the attentional benefit occurs at the intended saccade target rather than the actual landing position [7,34]. Accordingly, when the amplitude of the executed saccade is modified via **saccadic adaptation**, the metrics of the presaccadic shift of attention remain unaffected [38]; however, when the saccade target is constant throughout the experiment, the shift of attention can be linked to the adapted saccade endpoint [39,40]. Moreover, visual attention is not coupled to the endpoint of **global saccades**, which unintentionally land between two nearby potential saccade targets, but instead is allocated equally to both targets [41–43] (Figure 1C). Thus, whereas attentional orienting has often been considered to be dependent on oculomotor programming [25–29], converging behavioral evidence shows that presaccadic attention is tied to the intended motor goal rather than to the executed motor program [7,34,38,41–43], providing evidence for a dissociation of attentional and oculomotor orienting.



In addition, **antisaccade tasks** provide a unique opportunity to isolate and decouple the perceptual effects elicited by the visual stimulus and the final oculomotor command (Figure 1D). Visual spatial attention is simultaneously allocated to the cued location and the antisaccade goal ([44]; similar results but a different interpretation are given in [45]), suggesting attentional selection at both locations. If attention was functionally equivalent to saccade preparation [25], it should have been primarily allocated to the antisaccade goal. Following the same logic, deploying covert exogenous [33] or covert endogenous [31,32] attention does not speed up saccade initiation to the attended target, which would be expected if attention and oculomotor orienting were equivalent. In sum, these results show that covert attentional orienting is not the consequence of motor planning.

Modulations of visual representations by presaccadic attention and covert attention

The vast majority of presaccadic attention research has focused on its effects on performance (e.g., accuracy in discrimination tasks). We discuss here recent advances that go beyond mere performance measures and reveal how presaccadic attention modulates featural representations – in other words the processing of visual features, including orientation, spatial frequency, and contrast. This assessment enables a differentiation of the cognitive, neural, and computational processes underlying presaccadic and covert attention.

To characterize visual representation at the perceptual level, psychophysical procedures – for example, **adaptation**, **visual masking**, and **reverse correlation** – measure perceptual tuning functions. An adapter or a noise mask indexes how well human participants can detect or discriminate visual targets as a function of the distance between the target and the adaptor (or the mask) on a specific feature dimension (e.g., orientation, spatial frequency). The resulting perceptual tuning functions provide a gateway to compare perceptual effects resulting from different types of attention.

Orientation

Two studies employing psychophysical reverse correlation have investigated how presaccadic attention affects the perceptual orientation tuning function. In the first study [18], participants detected a visual target stimulus (a vertically oriented grating) presented at the saccade target. The target stimulus was embedded in random noise with rich orientation content, allowing an estimation of the perceptual orientation tuning function. Compared with a neutral condition, without saccadic eye movements, the orientation tuning function at the saccade target exhibited both gain enhancement (increased sensitivity) and tuning width narrowing (higher selectivity) (Figure 2A and Table 2). These modulations emerged ~100 ms before saccade onset and reached a maximum just before the eyes moved. Moreover, merely deploying covert attention within the same temporal interval without preparing a saccade did not alter performance. A follow-up study utilized dynamical noise and reported similar effects by presaccadic attention at the saccade target, and lower gain and a wider tuning function at a location opposite to the saccade direction, immediately before saccade onset [46].

Unlike presaccadic attention, both covert endogenous and exogenous spatial attention increase the gain of perceptual orientation (or motion) tuning functions without affecting its tuning width [47–51] (Figure 2B,C and Table 2). Only when participants are instructed to attend to a particular orientation, does covert feature-based attention reduce the width of orientation tuning [47,48].

Spatial frequency

Spatial frequency (SF) refers to the spatial scale (i.e., areas of relative light and dark in the visual scene) of visual inputs. The SF selectivity of the visual system varies across the visual field;





Figure 2. The effects of presaccadic and covert attention on featural representations. (A) In psychophysical experiments, participants were required to detect a grating at a fixed orientation (represented by 0° in the figure), and a reverse correlation technique was used to characterize the orientation tuning employed by the visual system. Presaccadic attention increases the gain and reduces the width of orientation tuning [18,46]. Panel adapted, with permission, from [18]. (B,C) Covert endogenous and exogenous attention only increase the gain [47–51]. Panels adapted, with permission, from [51]. (D–F) Presaccadic attention [18,54] and covert exogenous attention preferentially increase the sensitivity of high spatial frequency (SF) information by shifting the SF tuning curve rightward [51,55,56,61], whereas covert endogenous attention enhances a broad range of SFs uniformly [51,59–61]. The broken vertical lines indicate the peak of the tuning functions. Details of the experimental protocols can be found in Box 1. Panel (D) adapted, with permission, from [51]. (G–I) The subjective contrast appearance of visual stimuli can be estimated by measuring the point of subjective equality (PSE) in tasks requiring participants to compare the contrast of two stimuli. In the experiments, the contrast of a

(Figure legend continued at the bottom of the next page.)



sensitivity to higher SFs is higher at the fovea and decreases with eccentricity [1,2,52]. A study using reverse correlation revealed that presaccadic attention selectively increases sensitivity to higher SF information [18] (Figure 2D and Table 2). In this study, participants were required to detect a visual target stimulus (a vertically oriented grating embedded in random noise) presented at the saccade target. Compared with a neutral condition without saccadic eye movements, presaccadic attention increased the gain for higher SFs while leaving lower SFs unchanged, resulting in a rightward shift of the SF tuning function. Merely deploying covert attention within the same temporal interval without preparing a saccade did not alter performance. A follow-up study has confirmed the shift of SF tuning toward higher SF with presaccadic attention [53]. Moreover, this modulation is compulsory [54] and cannot be explained by a shift of the tuning function toward the SF of the target [18,54]: Using a visual masking procedure, it has been shown that the rightward shift of SF tuning by presaccadic attention could move the SF tuning function away from (higher than) the target, leading to an impairment on the task at hand (discriminating the orientation of the target) ([54]; details in Box 1).

Interestingly, the compulsory increment of high-SF sensitivity by presaccadic attention is reminiscent of the modulations by covert exogenous attention. In texture segmentation tasks (Box 1), exogenous attention increases spatial resolution, mediated by high SF, even when it impairs performance [52,55–58]. By contrast, covert endogenous attention always improves performance in the same tasks, without preferentially enhancing high-SF information [59,60] (Figure 2E,F and Table 2).

Converging evidence indicates that covert exogenous attention and endogenous attention modulate SF processing differently. In orientation discrimination tasks, regardless of stimulus eccentricity, exogenous attention preferentially enhances SFs higher than the peak frequency in the baseline condition, whereas endogenous attention enhances a broad range of lower and higher SFs [61]. Likewise, reverse correlation revealed that exogenous attention shifts peak sensitivity to higher SFs by enhancing the gain of SFs higher than the target SF, whereas endogenous attention enhances the gain of SFs below and above the SF of the target [51]. These changes in sensory tuning could underlie the differential performance effects of endogenous and exogenous attention in contrast sensitivity and texture segmentation tasks.

Contrast appearance

Covert attention changes the appearance of visual stimuli – how they appear subjectively to the participants – in a several static (e.g., contrast, SF, size) and dynamic (e.g., flicker, motion coherence) visual dimensions [62]. Does presaccadic attention similarly affect appearance?

The appearance of visual stimuli can be assessed by a comparative procedure in which experimenters measure the **point of subjective equality (PSE)** of a standard stimulus. Both exogenous [63–70] and endogenous [71] covert spatial attention increase perceived contrast of the stimulus at the attended location, as indicated by a shift of the PSE to a lower contrast (Figure 2H,I and Table 2). A counterpart of this effect has been reported for presaccadic attention (Table 2). As time approaches saccade onset, perceived contrast and orientation discrimination increase concurrently and progressively (Figure 2G). A control experiment with a cue-tostimulus interval matched to the presaccadic experiment demonstrated that endogenous

test stimulus typically varies across trials while the contrast of a standard stimulus is fixed. The percentage of trials in which participants judge the test stimulus to have a higher contrast than the standard (*y* axis) is plotted against the contrast of the test stimulus (*x* axis). The orange curves represent the condition in which the test stimulus is cued (attended). Both presaccadic attention [15] as well as endogenous [71] and exogenous [63–68] covert attention enhance perceived contrast. The enhancement by presaccadic attention exhibits a gradual trend right before saccade onset [the inset in panel (G) is adapted from [15]]. Panel (H) adapted from [62]; panel (I) adapted from [63].



Table 2. St	udies and experimental	procedures investigating	the modulation of visual	representations by	different types of attention in humans ^a

Feature	Method/task	Presaccadic attention	Covert endogenous attention	Covert exogenous attention
Orientation gain	Reverse correlation	 ✓ Li <i>et al.</i> 2016 [18] ✓ Ohl <i>et al.</i> 2017 [46] 	 ✓ Paltoglou and Neri 2012 [48] ✓ Wyart <i>et al.</i> 2012 [49] ✓ Fernández <i>et al.</i> 2021 [51] 	 ✓ Fernández <i>et al.</i> 2019 [50] ✓ Fernández <i>et al.</i> 2021 [51]
	Masking		✓ Baldassi and Verghese 2005 [47]	
Orientation tuning width	Reverse correlation	 ✓ Li <i>et al.</i> 2016 [18] ✓ Ohl <i>et al.</i> 2017 [46] 	 ✗ Paltoglou and Neri 2012 [48] ✗ Wyart <i>et al.</i> 2012 [49] ✗ Fernández <i>et al.</i> 2021 [51] 	 ✗ Fernández <i>et al.</i> 2019 [50] ✗ Fernández <i>et al.</i> 2021 [51]
	Masking		✗ Baldassi and Verghese 2005 [47]	
Enhancement of high spatial frequency	Texture segmentation		✗ Jigo and Carrasco 2018 [60]	 Yeshurun and Carrasco, 1998 [55] Yeshurun and Carrasco, 2000 [149] Talgar and Carrasco 2002 [58] Yeshurun and Carrasco 2008 [57] Jigo and Carrasco 2018 [60]
	Texture segmentation + adaptation		✗ Barbot and Carrasco 2017 [59]	✓ Carrasco et al. 2006 [56]
	Reverse correlation	✓ Li et al. 2016 [18]	X Fernández <i>et al.</i> 2021 [51]	 ✓ Fernández <i>et al.</i> 2021 [51] ✗ Fernández <i>et al.</i> 2019 [50]
	Masking	✔ Li et al 2019 [54]		X Talgar <i>et al.</i> 2000 [150]
	Orientation discrimination	✓ Kroell and Rolfs 2021 [53]	✗ Jigo and Carrasco 2020 [61]	✓ Jigo and Carrasco 2020 [61]
Contrast appearance	Comparative judgment (PSE) + orientation discrimination	✓ Rolfs and Carrasco 2012 [15]	✓ Liu <i>et al. 20</i> 09 [71]	 Carrasco et al. 2004 [63] Ling and Carrasco 2007 [64] Carrasco et al. 2008 [65] Störmer et al. 2009 [66] Anton-Erxleben et al. 2010 [67] Cutrone et al. 2014 [68] Barbot and Carrasco 2018 [69] Zhou et al. 2018 [70]
Contrast response function	Orientation discrimination	+ Li <i>et al.</i> 2021 [99]	* Ling and Carrasco 2006 [77] * Pestilli <i>et al.</i> 2009 [79] * or +: size-dependent Herrmann <i>et al.</i> 2010 [80] • Li <i>et al.</i> 2021 [99]	+ Ling and Carrasco 2006 [77] + Pestilli <i>et al.</i> 2007 [78] + Pestilli <i>et al.</i> 2009 [79] * or +: size-dependent Herrmann <i>et al.</i> 2010 [80] + Fernández and Carrasco 2020 [145] • Li <i>et al.</i> 2021 [99]
	Contrast discrimination		+ Morrone <i>et al.</i> 2002 [74] + Morrone <i>et al.</i> 2004 [75] ♦ Huang and Dobkins 2005 [76]	

^aKey: \checkmark , the feature modulation indicated in the first column is observed; \bigstar , the feature modulation indicated in the first column is not observed; *, contrast gain; +, response gain; +, mixture of both contrast gain and response gain.

attention cannot modulate contrast appearance in such a rapid fashion [15]. Thus, even though the perceptual modulations of contrast appearance are similar for presaccadic attention and covert endogenous attention, their temporal dynamics differ. These results link the dynamics of saccade preparation, visual performance, and subjective experience, and show that upcoming eye movements alter visual processing by increasing signal strength.

Contrast response functions and computational models

Computational models of attention aim to characterize attentional modulation of the inputoutput functions of visual neurons (e.g., contrast response functions: neural responses as a function of stimulus contrast). The neural activity as a function of stimulus contrast is either



Box 1. Texture segmentation

Texture segmentation refers to the process by which the visual system isolates a region from its background based on its local structure. In the example depicted in Figure I, participants were required to detect the presence of the target patch whose line elements differ in orientation from the otherwise uniform background. Performance exhibits an inverted U-shape as a function of target eccentricity: higher performance for the target at midperiphery than at central and peripheral locations. This central performance drop (CPD) indicates that the resolution of the visual system, or sensitivity to high spatial frequency (SF) information, is too high for target detection. Accordingly, peak performance shifts to more peripheral or central locations when the texture scale increases or decreases, respectively [55]. Selective adaptation to high-SF filters reduces participants' sensitivity to high SF, and correspondingly the CPD [56,59].

Compared with a neutral baseline condition, exogenous attention impairs performance when the target patch is near the fovea, where visual resolution is already too high for the task, but improves performance when the target patch is at periphery [55,56,58,60,149]. These results reveal that exogenous attention increases visual resolution, or the sensitivity to high SF, in a compulsory manner, even when detrimental for the task at hand. In the same task, endogenous attention adjusts resolution and improves performance regardless of target eccentricity [59,60]. Critically, selective adaptation to high SF eliminates the central attentional impairment observed with exogenous attention [56] as well as the central improvement observed with endogenous attention [59].

Visual masking was utilized to test the effect of presaccadic attention on the processing of SF [54] (Figure II). A target at a fixed SF was superimposed with noise (mask) of different (lower, same, or higher) SFs. In the neutral baseline condition, discrimination of the target orientation was lowest when the target was paired with the noise containing the same SF. Presaccadic attention enhanced the discriminability of the target when the noise had a lower or the same SF as the target, but impaired its discriminability when the SF of the noise was higher than that of the target, presumably because it increased the suppressive effect elicited by the high-SF noise. Merely deploying covert endogenous attention within the same temporal interval did not alter performance. These results indicate that presaccadic attention enhances high SF information, even when detrimental for task performance.



Figure I. Texture segmentation tasks. Top: example stimuli. Bottom: exogenous attention impairs or improves performance as a function of eccentricity; endogenous attention improves performance throughout eccentricity. Adapted from [52].









measured directly or is simulated by computational models and fit to behavioral performance. For covert attention, mainly two types of attentional modulation have been reported in both neurophysiological [72,73] and behavioral studies [74–80]: a contrast gain change, in which the response function shifts horizontally, as if attention scales the input stimulus contrast (e.g., Figure 3A), and a response gain change, in which the asymptotic response moves vertically, as if attention multiplicatively scales the output response of the neurons (Figure 3B). In addition, human functional magnetic resonance imaging (fMRI) studies have reported an additive shift of response functions [81,82], or a combination of additive shift and contrast gain change by attention [83].

Various computational studies have modeled attentional modulation of neural responses [84-87]. However, computational models of visual attention have either referred to covert attention or presaccadic attention exclusively, or have referred to spatial covert attention as a unitary construct. We focus here on a class of models that rely on normalization [88] to describe the computations underlying visual attention [72,73,89–91] because these normalization models of attention (NMA) have been widely applied to explain both behavioral [80,91,92] and neurophysiological [93-97] data in recent years. In normalization, the response of a visual neuron is modeled by the excitatory drive divided by the suppressive drive (normalization pool), computed as the excitatory drive summed over a pool of neurons [88]. In the NMA of Reynolds and Heeger [73], attention is modeled by attentional gain factors that multiplicatively modulate the excitatory drive of the neurons before normalization (Figure 3C). This computation predicts that attention exhibits a response gain (contrast gain) change when the size of the attentional field is small (large) relative to the stimulus size (Figure 3A,B). An intermediate attentional field size will lead to a mix of both response gain and contrast gain changes. Behavioral studies on covert attention have confirmed this prediction [80,92]. The additive shift by attention observed in human fMRI can also be explained by applying the NMA to a neural population consisting of neurons with heterogeneous (position and orientation) tuning, and summing activity across many neurons to simulate the response of fMRI voxels [98].

Interestingly, unlike covert attention, presaccadic attention generates response gain changes regardless of the size of the attentional field (Figure 3A,C and Table 2) [99]. Model comparisons revealed that, whereas modulations by covert attention are best fit by the NMA, modulations by presaccadic attention are best fit by a response gain model in which response gain factors





Trends in Neurosciences

Figure 3. Attentional modulations of contrast response functions. (A) Illustrations of how attentional field size (relative to the stimulus size) affects the form of attentional modulation. The left column represents the condition in which the size of the attentional field is large, whereas the right column represents the condition in which the size of the attentional field is small. Covert attention (endogenous and exogenous; top row) generates contrast gain when the size of the attentional field is large and response gain when the size of the attentional field is small [73,80]. By contrast, presaccadic attention (bottom row) generates response gain changes regardless of the attentional field size [99]. The insets in the bottom row illustrate how the size of attention field was manipulated by the location uncertainty of the target in a study of presaccadic attention [99]. Only the fixation point with the pre-cue and the right half of the screen are shown. The left half of the screen contains similar stimuli (an aperture outlined by four black dots and a test stimulus). In the experiments, participants made a saccadic eye movement toward the center of the cued aperture (outlined by the four black dots). In the large attention field size condition, the aperture was large and the test stimulus (a grating) was presented at one of five possible locations (indicated by green circles for illustration purposes only) with equal probability. In the small attention field size condition, the test stimulus was presented at a fixed location [73,80,99] (attentional field size was also manipulated by location uncertainty in covert attention experiments [73,80]). (B) The Reynolds and Heeger normalization model of attention (NMA) explains these size-dependent gain changes [73]. Blue-yellow 2D images represent a population of visual neurons selective for different orientations and receptive field (RF) centers. The response of visual neurons is computed by normalization: dividing the excitatory drive of neurons by their suppressive drive. The excitatory drive is determined by the preferred orientation and position of each neuron. The suppressive drive is computed by convolving the excitatory drive with the suppression kernel. The suppression kernel is uniform in the orientation domain. For simulating surround suppression of visual neurons, the suppression kernel is a Gaussian with a wide width in its spatial domain. In the NMA, attention is modeled as attentional gain factors that multiplicatively modulate the excitatory drive before computing the suppressive drive and normalization. (C) The modulation of presaccadic attention is better explained by response gain factors that scale the responses of neurons after normalization. In both (B) and (C) it is assumed that attention is deployed to the center of the target stimulus (the vertical grating). Panels (A-C) adapted, with permission, from [99].

modulate the response of the neurons after normalization (Figure 3C). These results indicate that the computations underlying presaccadic attention and covert attention are different [99].

Neural correlates of presaccadic and covert attention

Common brain areas but distinct subpopulations

The tight coupling between eye movements and attentional orienting raises the question of whether presaccadic and covert spatial attention are based on the same neural processes. At a broad scale, the brain structures that are active during saccadic eye movements [the **frontal** eye fields (FEF), the precentral sulcus, and the **superior colliculus (SC)**] are also selectively modulated during covert attention tasks in human- and non-human primates [100–103].

CellPress

Feedback signals projecting from these areas to early visual cortices are assumed to enhance visual processing [104,105] and thereby account for the commonly observed presaccadic attentional effects.

In non-human primate studies, subthreshold microstimulation of FEF and SC – which would elicit a saccade if stimulated above threshold – modulates activity in visual cortex [106] and enhances visual sensitivity at the movement field of the stimulated neurons (Figure 1E) [107,108]. However, FEF and SC have distinct neuronal populations that are related to visual and motor activity [108–113]. Because microstimulation modulates both visual and motor cells, it cannot distinguish between oculomotor and attentional orienting, which are controlled by distinct populations within these overlapping neuronal circuits: Visual and visuomotor cells exhibit sustained activity during covert attentional orienting, but motor cells remain silent [110,111] and have disproportionately little feedback connections to early visual cortex [112]. Furthermore, covertly attended locations and eye movement goals are represented by different subsets of spatially tuned neurons in frontal cortex, contradicting the idea that the control of spatial attention is dependent on oculomotor control circuits [113].

Likewise, in humans, the early laterality of visual evoked event-related potentials (ERPs) correlates with covert but not presaccadic attention [114], and transcranial magnetic stimulation (TMS) over FEF has revealed a clear temporal dissociation of saccade preparation and attention [115]. These results are consistent with the findings that saccade preparation and covert attention – despite involving overlapping neural regions – are neurophysiologically distinct phenomena [109,116]. In summary, neurophysiological evidence reveals that distinct neural populations in FEF and SC control covert attention and saccade preparation in human and non-human primates, potentially giving rise to differences in their perceptual correlates.

Response enhancement and noise reduction

Both endogenous and exogenous attention enhance neural response amplitude at the attended location throughout the visual system [72,117]. Response enhancement by presaccadic attention has also been observed in two macaque brain regions – **visual area V1** [118] and **visual area V4** [119,120] – and throughout both ventral and dorsal visual streams in human cortex [121]. The enhanced response is accompanied by an increment of decodable stimulus information (orientation) from the neural population just before saccade onset [122], matching the presaccadic attentional effects in behavioral studies. Likewise, in line with behavioral evidence (e.g., [44]), response enhancement by covert attention and presaccadic attention coexist, as revealed by an antisaccade experiment reporting qualitatively similar activity modulations corresponding to an endogenously attended location and the opposite saccade target location [123].

Recording in V4 has also revealed that both covert attention [124–126] and presaccadic attention [127] reduce the variability and noise correlation among visual neurons, which correlate with changes in behavioral performance [128].

Featural representation

How do presaccadic and covert attention modulate featural representations at the neural level? In line with psychophysical findings [47–49], endogenous spatial attention increases the gain (amplitude) on orientation tuning functions of visual neurons without affecting their tuning width [129,130]. To our knowledge, there is no parallel neurophysiological study for exogenous attention. Studies in V4 have shown that presaccadic attention enhances the response of visual neurons more to the



preferred orientation than to nonpreferred orientations [119] and increases the gain on orientation tuning functions, but did not report whether tuning width was affected [123]. Thus, whereas endogenous attention does not affect neural orientation tuning width, it remains unknown whether presaccadic attention does, because its effects on gain and tuning have not been assessed independently.

Two (nonmutually exclusive) neural mechanisms may underlie the increment of sensitivity to high-SF information by spatial attention: first, receptive field (RF) shifts: the deployment of endogenous attention can shift visual neuron RF toward the attended location. In monkey single-unit recording, this RF shift occurs in both ventral [131] and dorsal extrastriate visual cortex [132]. Likewise, human neuroimaging studies measuring population receptive fields (pRFs) reported a shift of pRFs toward the attended location in the visual cortex [133]. The RF shifts enhance the neural representation of the attended stimuli and are accompanied by a shrinkage of RF size at the attended location [132]. In principle, smaller RF sizes could benefit the discrimination of smaller targets in texture segmentation tasks because a smaller RF would reduce the interference from background stimuli [2]. Similarly, just before monkeys make saccadic eye movements, RFs shift toward the saccade target in response to presaccadic visual stimuli in V4 [120,134] and prefrontal cortex [135]. In sum, RF shifts have been observed for endogenous and presaccadic attention, but comparable studies for exogenous attention are lacking. However, at the behavioral level, only presaccadic attention and exogenous attention shift behavioral SF tuning toward higher SFs (Figure 2 and Table 2). Endogenous attention conceivably evokes additional mechanisms that benefit the processing of low-SF information, thus leaving the overall SF sensitivity tuning function unchanged.

The second potential neural mechanism to support an increment in high-SF sensitivity is a preferential enhancement of visual neurons tuned to high SFs. Neurons in the mouse primary visual cortex exhibit gain increments during a heightened attentional state, as behaviorally assessed by locomotion and a dilated pupil. Critically, neurons selective for higher SFs exhibit larger gain increments than those selective for lower frequencies [136]. In mice, the dorsal lateral geniculate nucleus of the thalamus also shows a preferential gain increment in locomotion for neurons selective for high SF [137]. These neural modulations could support the perceptual effects observed in behavioral attention research, but neurophysiological studies using a spatial attention protocol have yet to test this assumption. A critical distinction is that the effect of locomotion is global across the entire visual field whereas the effect of spatial attention is restricted to the attended location. In general, these observations are consistent with psychophysical experiments employing a selective adaptation protocol, which revealed that spatial attention tunes SF sensitivity via high-SF channels [56,59].

Concluding remarks

Although both presaccadic attention and covert spatial attention enable selective processing of visual information and facilitate perception, there are clear dissociations in their temporal dynamics, modulations of featural representation of basic visual dimensions, neural computations, and neural correlates. Although questions remain about the specific origin, nature, and functional significance of presaccadic attention and covert attention, converging evidence indicates that the notion that visual attention cannot be decoupled from eye movement programming should be revised. Presaccadic and covert attention are distinct neurocognitive processes that do not depend on one another.

The dissociations between covert attention and presaccadic attention provide important insight into brain function and call for further investigation. Our understanding of presaccadic attention –

Outstanding questions

Do presaccadic attentional modulations of featural representations, as suggested, support trans-saccadic integration and visual stability at around the time of saccades?

Behaviorally, when participants plan a saccade, presaccadic attention enhances gain, narrows orientation tuning, and shifts SF selectivity at the saccade target. Is this also the case when participants saccade in response to a salient stimulus, without voluntarily planning to do so?

Presaccadic attention changes contrast appearance. Does it also change the appearance of other static (e.g., spatial frequency, size) and dynamic (e.g., flicker, speed) visual dimensions, as covert attention does?

Presaccadic attention alters featural representations at the saccade target. What happens simultaneously to featural representations at other locations, for example, at the fovea (where the gaze is about to leave) and at locations around the saccade target?

Presaccadic attention affects encoding of sensory information. Does it also affect other processes such as information readout or decision criteria?

Presaccadic attention alters featural representations at the saccade target. What neural correlates of these effects would neurophysiological and neuroimaging studies reveal?

Reaction time studies with special populations (e.g., Moebius syndrome) inform our knowledge of eye movement programming/execution and covert attention. Would discrimination tasks yield consistent findings? Would people with such oculomotor conditions show the same modulations of featural representations as people with intact oculomotor systems?

TMS has shown a causal role of occipital cortex in exogenous attention. Does the occipital cortex also play a causal role in presaccadic attention, and how does it interact with higher brain areas (e.g., FEF)?



as well as of its similarities to and differences from covert spatial attention – would be further advanced by integrating knowledge gathered from psychophysics, neuropsychology, neurophysiology, neuroimaging, and computational models in human and non-human primates, while keeping the experimental protocols as similar as possible (see Outstanding questions). Moreover, it would be informative to isolate individual components of eye movements (planning, programming, and execution), which may differentially influence the findings in studies with human participants and non-human primates. Depending on the time-period of the analysis, different oculomotor components could contribute to the effects of presaccadic attention or conflate the effects attributed to covert attention.

The findings that covert and presaccadic attention lead to different modulations of featural representations suggest that they serve different functional roles. We have proposed that the sharpening of orientation tuning [18], enhancement of high SF [18,54], increase in perceived contrast [15], and the overall amplification of the contrast response [99] elicited by presaccadic attention may contribute to **trans-saccadic integration**, serving perceptual continuity across saccades by making the peripheral saccade target more fovea-like before it is shifted to the fovea. Neurophysiologists have also suggested that the orientation gain increment observed in monkey visual cortex could aid trans-saccadic integration [119]. This notion that presaccadic attention may contribute to trans-saccadic integration is rooted in the idea of Bayesian integration that the sameness of two inputs makes them more likely to be considered to have a common source [138,139].

This review has focused primarily on the differences among presaccadic, covert exogenous, and covert endogenous attention regarding their effects on sensory encoding, but their differences may also manifest in other processes. There is evidence, for instance, that covert endogenous attention also affects information processing after sensory encoding because neural activity is also selectively modulated during readout of visual information in occipital cortex, TPJ (temporoparietal junction), and precentral sulcus/FEF [140–144]. Likewise, it remains unknown whether presaccadic attention modulates neural activity during readout and, if so, whether such modulation differs from that of exogenous or endogenous covert attention. To further our understanding of presaccadic attention and to compare it with endogenous and exogenous covert attention, it would be informative to conduct analogous studies assessing the modulatory effects of presaccadic attention on neural activity during encoding and readout.

In addition, TMS has been successfully used in human participants to identify brain regions causally related to visual attention. For example, by briefly disrupting cortical excitability of the occipital cortex with TMS the effects of covert exogenous attention at attended and unattended locations are extinguished [145]. Likewise, TMS of specific areas of the dorsal (rIPS, intraparietal sulcus) and ventral (rTPJ) attention networks abolishes benefits from exogenous cueing, showing that both networks are implicated in exogenous attention [146]. Furthermore, TMS of FEF has been shown to affect presaccadic attention [147]. Future studies are necessary to test the effects of presaccadic and covert attention in multiple brain regions to allow systematic comparisons of the necessary roles of specific brain regions and neural pathways in humans.

The perceptual consequences, featural representations, and neural computations and correlates of covert attention have been well characterized, but more so for the endogenous than the exogenous type (reviewed in [1,2,52,62,72,117]). Likewise, the characterization of presaccadic attention has concentrated more on tasks in which participants prepare a saccade to a target

How would to-be-developed models encompassing both covert and presaccadic attention advance our knowledge of visual perception by taking into account the dissociations reviewed here?

location voluntarily (according to a central cue) than involuntarily (reflexively saccading to a peripheral target). Further investigating exogenous attention and involuntary eye movements, as well as comparing their perceptual consequences, featural representations, computations, and neural correlates with endogenous attention and voluntary eye movements, will advance our knowledge of how presaccadic attention and covert attention modulate visual selection and perception.

Acknowledgments

This research was supported by National Institutes of Health National Eye Institute grant R01 EY019693 to M.C. and a Feodor Lynen Research Fellowship from the Alexander von Humboldt Foundation to N.M.H. We thank Antoine Barbot, Antonio Fernández, Marc Himmelberg, Michael Jigo, and other members of the laboratory of M.C., as well as Luca Wollenberg and Heiner Deubel, for useful comments.

Declaration of interests

The authors declare no competing interests.

References

- 1. Carrasco, M. (2011) Visual attention: the past 25 years. *Vis. Res.* 51, 1484–1525
- Anton-Erxleben, K. and Carrasco, M. (2013) Attentional enhancement of spatial resolution: linking behavioural and neurophysiological evidence. *Nat. Rev. Neurosci.* 14, 188–200
- Binda, P. and Morrone, M.C. (2018) Vision during saccadic eye movements. Annu. Rev. Vis. Sci. 4, 193–213
- Marino, A.C. and Mazer, J.A. (2016) Perisaccadic updating of visual representations and attentional states: linking behavior and neurophysiology. *Front. Syst. Neurosci.* 10, 3
- Kowler, E. et al. (1995) The role of attention in the programming of saccades. Vis. Res. 35, 1897–1916
- Hoffman, J.E. and Subramaniam, B. (1995) The role of visual attention in saccadic eye movements. *Percept. Psychophys.* 57, 787–795
- Deubel, H. and Schneider, W.X. (1996) Saccade target selection and object recognition: evidence for a common attentional mechanism. Vis. Res. 36, 1827–1837
- 8. Hanning, N.M. *et al.* (2019) Sensitivity measures of visuospatial attention. *J. Vis.* 19, 17
- Ghahghaei, S. and Verghese, P. (2017) Texture segmentation influences the spatial profile of presaccadic attention. J. Vis. 17, 10
- 10. Puntiroli, M. *et al.* (2018) Placeholder objects shape spatial attention effects before eye movements. *J. Vis.* 18, 1
- Szinte, M. et al. (2019) The spread of presaccadic attention depends on the spatial configuration of the visual scene. Sci. Rep. 9, 14034
- 12. Castet, E. *et al.* (2006) Dynamics of attentional deployment during saccadic programming. *J. Vis.* 6, 196–212
- Deubel, H. (2008) The time course of presaccadic attention shifts. *Psychol. Res.* 72, 630–640
 Filali-Sadouk, N. *et al.* (2010) Similar effect of cueing conditions
- Filali-Sadouk, N. *et al.* (2010) Similar effect of cueing conditions on attentional and saccadic temporal dynamics. *J. Vis.* 10, 21
 Rolfs, M. and Carrasco, M. (2012) Rapid simultaneous en-
- Hons, M. and Garasso, M. (2012) hapd similar leads of r hancement of visual sensitivity and perceived contrast during saccade preparation. J. Neurosci. 32, 13744–13752
- 16. Harrison, W.J. *et al.* (2013) Eye movement targets are released from visual crowding. *J. Neurosci.* 33, 2927–2933
- White, A.L. *et al.* (2013) Adaptive deployment of spatial and feature-based attention before saccades. *Vis. Res.* 85, 26–35
- Li, H.-H. et al. (2016) Saccade preparation reshapes sensory tuning. Curr. Biol. 26, 1564–1570
- Jonikaitis, D. et al. (2017) Spatial attention during saccade decisions. J. Neurophysiol. 118, 149–160
- Hunt, A.R. et al. (2019) The relationship between spatial attention and eye movements. Curr. Top. Behav. Neurosci. 41, 255–278
- 21. Montagnini, A. and Castet, E. (2007) Spatiotemporal dynamics of visual attention during saccade preparation: Independence

and coupling between attention and movement planning. J. Vis. 7, 8

- Zhao, M. et al. (2012) Eye movements and attention: the role of pre-saccadic shifts of attention in perception, memory and the control of saccades. Vis. Res. 74, 40–60
- Khan, A.Z. et al. (2015) Saccade execution suppresses discrimination at distractor locations rather than enhancing the saccade goal location. Eur. J. Neurosci. 41, 1624–1634
- Kreyenmeier, P. et al. (2020) Theory of visual attention (TVA) in action: assessing premotor attention in simultaneous eye-hand movements. *Cortex.* 133, 133–148
- Rizzolatti, G. et al. (1987) Reorienting attention across the horizontal and vertical meridians: evidence in favor of a premotor theory of attention. *Neuropsychologia*. 25, 31–40
- Craighero, L. et al. (2001) Peripheral oculomotor palsy affects orienting of visuospatial attention. Neuroreport 12, 3283–3286
- Smith, D.T. et al. (2004) Exogenous orienting of attention depends upon the ability to execute eye movements. *Curr. Biol.* 14, 792–795
- Smith, D.T. *et al.* (2010) Deficits of reflexive attention induced by abduction of the eye. *Neuropsychologia* 48, 1269–1276
- Smith, D.T. et al. (2012) Saccade preparation is required for exogenous attention but not endogenous attention or IOR. J. Exp. Psychol. Hum. Percept. Perform. 38, 1438–1447
- Casteau, S. and Smith, D.T. (2020) Covert attention beyond the range of eye-movements: Evidence for a dissociation between exogenous and endogenous orienting. *Cortex* 122, 170–186
- Hunt, A.R. and Kingstone, A. (2003) Covert and overt voluntary attention: linked or independent? *Brain Res. Cogn. Brain Res.* 18, 102–105
- Belopolsky, A.V. and Theeuwes, J. (2009) When are attention and saccade preparation dissociated? *Psychol. Sci.* 20, 1340–1347
- MacLean, G.H. et al. (2015) Does oculomotor readiness mediate exogenous capture of visual attention? J. Exp. Psychol. Hum. Percept. Perform. 41, 1260–1270
- Hanning, N.M. et al. (2019) Visual attention is not limited to the oculomotor range. Proc. Natl. Acad. Sci. 116, 9665–9670
- Hanning, N.M. and Deubel, H. (2020) Attention capture outside the oculomotor range. *Curr. Biol.* 30, R1353–R1355
- Masson, N. et al. (2020) Exogenous covert shift of attention without the ability to plan eye movements. *Curr. Biol.* 30, R1032–R1033
- Carrasco, M. and Hanning, N.M. (2020) Visual perception: attending beyond the eyes' reach. *Curr. Biol.* 30, R1322–R1324
- Ditterich, J. et al. (2000) Relation between the metrics of the presaccadic attention shift and of the saccade before and after saccadic adaptation. J. Neurophysiol. 84, 1809–1813
- Doré-Mazars, K. and Collins, T. (2005) Saccadic adaptation shifts the pre-saccadic attention focus. *Exp. Brain Res.* 162, 537–542



CellPress

Trends in Neurosciences

- Collins, T. et al. (2010) Visual target selection and motor planning define attentional enhancement at perceptual processing stages. Front. Hum. Neurosci. 4, 14
- Van der Stigchel, S. and de Vries, J.P. (2015) There is no attentional global effect: attentional shifts are independent of the saccade endpoint. J. Vis. 15, 17
- Wollenberg, L. et al. (2018) Visual attention is not deployed at the endpoint of averaging saccades. PLoS Biol. 16, e2006548
- Wollenberg, L. et al. (2020) Visual attention and eye movement control during oculomotor competition. J. Vis. 20, 16
- 44. Klapetek, A. *et al.* (2016) Attention allocation before antisaccades. *J. Vis.* 16, 11
- Mikula, L. *et al.* (2018) Spatial and temporal dynamics of presaccadic attentional facilitation before pro- and antisaccades. *J. Vis.* 18, 2
- 46. Ohl, S. et al. (2017) Selective enhancement of orientation tuning before saccades. J. Vis. 17, 2
- Baldassi, S. and Verghese, P. (2005) Attention to locations and features: different top-down modulation of detector weights. *J. Vis.* 5, 556–570
- Paltoglou, A.E. and Neri, P. (2012) Attentional control of sensory tuning in human visual perception. J. Neurophysiol. 107, 1260–1274
- Wyart, V. et al. (2012) Dissociable prior influences of signal probability and relevance on visual contrast sensitivity. Proc. Natl. Acad. Sci. U. S. A. 109, 3593–3598
- 50. Fernández, A. *et al.* (2019) How exogenous spatial attention affects visual representation. *J. Vis.* 19, 4
- Fernández, A. et al. (2021) Differential effects of endogenous and exogenous attention on sensory tuning. *BioRxiv* Published online April 4, 2021. https://doi.org/10.1101/2021.04.03.438325
- Carrasco, M. and Barbot, A. (2014) How attention affects spatial resolution. *Cold Spring Harb. Symp. Quant. Biol.* 79, 149–160
- Kroell, L.M. and Rolfs, M. (2021) The peripheral sensitivity profile at the saccade target reshapes during saccade preparation. *Cortex* 139, 12–26
- Li, H.-H. et al. (2019) Presaccadic attention improves or impairs performance by enhancing sensitivity to higher spatial frequencies. Sci. Rep. 9, 2659
- Yeshurun, Y. and Carrasco, M. (1998) Attention improves or impairs visual performance by enhancing spatial resolution. *Nature*. 396, 72–75
- Carrasco, M. et al. (2006) How attention enhances spatial resolution: evidence from selective adaptation to spatial frequency. Percept. Psychophys. 68, 1004–1012
- Yeshurun, Y. and Carrasco, M. (2008) The effects of transient attention on spatial resolution and the size of the attentional cue. *Percept. Psychophys.* 70, 104–113
- Talgar, C.P. and Carrasco, M. (2002) Vertical meridian asymmetry in spatial resolution: visual and attentional factors. *Psychon. Bull. Rev.* 9, 714–722
- Barbot, A. and Carrasco, M. (2017) Attention modifies spatial resolution according to task demands. *Psychol. Sci.* 28, 285–296
- Jigo, M. and Carrasco, M. (2018) Attention alters spatial resolution by modulating second-order processing. J. Vis. 18, 2
- Jigo, M. and Carrasco, M. (2020) Differential impact of exogenous and endogenous attention on the contrast sensitivity function across eccentricity. J. Vis. 20, 11
- 62. Carrasco, M. and Barbot, A. (2019) Spatial attention alters visual appearance. *Curr. Opin. Psychol.* 29, 56–64
- 63. Carrasco, M. *et al.* (2004) Attention alters appearance. *Nat. Neurosci.* 7, 308–313
- Ling, S. and Carrasco, M. (2007) Transient covert attention does alter appearance: a reply to Schneider (2006). *Percept. Psychophys.* 69, 1051–1058
- Carrasco, M. et al. (2008) Transient attention does increase perceived contrast of suprathreshold stimuli: a reply to Prinzmetal, Long, and Leonhardt (2008). Percept. Psychophys. 70, 1151–1164
- Störmer, V.S. et al. (2009) Cross-modal cueing of attention alters appearance and early cortical processing of visual stimuli. Proc. Natl. Acad. Sci. U. S. A. 106, 22456–22461

- Anton-Erxleben, K. *et al.* (2010) Evaluating comparative and equality judgments in contrast perception: attention alters appearance. *J. Vis.* 10, 6
- Cutrone, E.K. *et al.* (2014) Attention enhances contrast appearance via increased input baseline of neural responses. *J. Vis.* 14, 16
- Barbot, A. and Carrasco, M. (2018) Emotion and anxiety potentiate the way attention alters visual appearance. *Sci. Rep.* 8, 5938
- Zhou, L.-F. *et al.* (2018) Attentional effect on contrast appearance: from enhancement to attenuation. *J. Exp. Psychol. Hum. Percept. Perform.* 44, 806–817
- Liu, T. et al. (2009) Voluntary attention enhances contrast appearance. Psychol. Sci. 20, 354–362
- Maunsell, J.H.R. (2015) Neuronal mechanisms of visual attention. Annu. Rev. Vis. Sci. 1, 373–391
- 73. Reynolds, J.H. and Heeger, D.J. (2009) The normalization model of attention. *Neuron.* 61, 168–185
- Morrone, M.C. et al. (2002) Color and luminance contrasts attract independent attention. Curr. Biol. 12, 1134–1137
- Morrone, M.C. et al. (2004) Different attentional resources modulate the gain mechanisms for color and luminance contrast. Vis. Res. 44, 1389–1401
- Huang, L. and Dobkins, K.R. (2005) Attentional effects on contrast discrimination in humans: evidence for both contrast gain and response gain. *Vis. Res.* 45, 1201–1212
- Ling, S. and Carrasco, M. (2006) Sustained and transient covert attention enhance the signal via different contrast response functions. *Vis. Res.* 46, 1210–1220
- Pestilli, F. et al. (2007) How do attention and adaptation affect contrast sensitivity? J. Vis. 7, 9
- Pestilii, F. et al. (2009) A population-coding model of attention's influence on contrast response: estimating neural effects from psychophysical data. Vis. Res. 49, 1144–1153
- Hermann, K. *et al.* (2010) When size matters: attention affects performance by contrast or response gain. *Nat. Neurosci.* 13, 1554–1559
- Buracas, G.T. and Boynton, G.M. (2007) The effect of spatial attention on contrast response functions in human visual cortex. J. Neurosci. 27, 93–97
- Murray, S.O. (2008) The effects of spatial attention in early human visual cortex are stimulus independent. J. Vis. 8, 2
- Li, X. et al. (2008) Blood oxygenation level-dependent contrast response functions identify mechanisms of covert attention in early visual areas. Proc. Natl. Acad. Sci. U. S. A. 105, 6202–6207
- Bundesen, C. *et al.* (2005) A neural theory of visual attention: bridging cognition and neurophysiology. *Psychol. Rev.* 112, 291–328
- Bhatt, R. *et al.* (2007) Texture segregation by visual cortex: perceptual grouping, attention, and learning. *Vis. Res.* 47, 3173–3211
- Womelsdorf, T. *et al.* (2008) Receptive field shift and shrinkage in macaque middle temporal area through attentional gain modulation. *J. Neurosci.* 28, 8934–8944
- Schneider, W.X. (2013) Selective visual processing across competition episodes: a theory of task-driven visual attention and working memory. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 368, 20130060
- Carandini, M. and Heeger, D.J. (2011) Normalization as a canonical neural computation. *Nat. Rev. Neurosci.* 13, 51–62
- Lee, J. and Maunsell, J.H.R. (2009) A normalization model of attentional modulation of single unit responses. *PLoS One* 4, e4651
- Boynton, G.M. (2009) A framework for describing the effects of attention on visual responses. *Vis. Res.* 49, 1129–1143
- Jigo, M. et al. (2021) How voluntary and involuntary attention differentially shape spatial resolution. *BioRxiv* Published online March 19, 2021. https://doi.org/10.1101/2021.01.26.428173
- Herrmann, K. et al. (2012) Feature-based attention enhances performance by increasing response gain. Vis. Res. 74, 10–20
- Ni, A.M. *et al.* (2012) Tuned normalization explains the size of attention modulations. *Neuron* 73, 803–813
- 94. Schwedhelm, P. et al. (2016) An extended normalization model of attention accounts for feature-based attentional enhancement



of both response and coherence gain. *PLoS Comput. Biol.* 12, e1005225

- Verhoef, B.-E. and Maunsell, J.H. (2016) Attention operates uniformly throughout the classical receptive field and the surround. *eLife* 5, e17256
- Verhoef, B.-E. and Maunsell, J.H.R. (2017) Attention-related changes in correlated neuronal activity arise from normalization mechanisms. *Nat. Neurosci.* 20, 969–977
- Ni, A.M. and Maunsell, J.H.R. (2019) Neuronal effects of spatial and feature attention differ due to normalization. *J. Neurosci.* 39, 5493–5505
- Hara, Y. et al. (2014) Differing effects of attention in single-units and populations are well predicted by heterogeneous tuning and the normalization model of attention. Front. Comput. Neurosci. 8, 12
- Li, H.-H. et al. (2021) Different computations underlie overt presaccadic and covert spatial attention. Nat. Hum. Behav. Published online April 19, 2021. https://doi.org/10.1038/ s41562-021-01099-4
- Nobre, A.C. et al. (2000) Covert visual spatial orienting and saccades: overlapping neural systems. Neuroimage 11, 210–216
- Bogadhi, A.R. et al. (2018) Brain regions modulated during covert visual attention in the macaque. Sci. Rep. 8, 15237
- Bollimunta, A. et al. (2018) Comparing frontal eye field and superior colliculus contributions to covert spatial attention. Nat. Commun. 9, 3553
- 103. Jonikaitis, D. and Moore, T. (2019) The interdependence of attention, working memory and gaze control: behavior and neural circuitry. *Curr. Opin. Psychol.* 29, 126–134
- Ekstrom, L.B. *et al.* (2008) Bottom-up dependent gating of frontal signals in early visual cortex. *Science* 321, 414–417
- Bisley, J.W. and Mirpour, K. (2019) The neural instantiation of a priority map. *Curr. Opin. Psychol.* 29, 108–112
- Moore, T. and Armstrong, K.M. (2003) Selective gating of visual signals by microstimulation of frontal cortex. *Nature* 421, 370–373
- 107. Moore, T. and Fallah, M. (2004) Microstimulation of the frontal eye field and its effects on covert spatial attention. *J. Neurophysiol.* 91, 152–162
- Müller, J.R. et al. (2005) Microstimulation of the superior colliculus focuses attention without moving the eyes. Proc. Natl. Acad. Sci. U. S. A. 102, 524–529
- 109. Juan, C.-H. et al. (2004) Dissociation of spatial attention and saccade preparation. Proc. Natl. Acad. Sci. U. S. A. 101, 15541–15544
- Ignashchenkova, A. et al. (2004) Neuron-specific contribution of the superior colliculus to overt and covert shifts of attention. *Nat. Neurosci.* 7, 56–64
- Gregoriou, G.G. *et al.* (2012) Cell-type-specific synchronization of neural activity in FEF with V4 during attention. *Neuron* 73, 581–594
- 112. Merrikhi, Y. et al. (2017) Spatial working memory alters the efficacy of input to visual cortex. Nat. Commun. 8, 15041
- 113. Messinger, A. (2021) Separable neuronal contributions to covertly attended locations and movement goals in macaque frontal cortex. *Sci. Adv.* 7, eabe0716
- 114. Hilo-Merkovich, R. et al. (2018) Task performance in covert, but not overt, attention correlates with early laterality of visual evoked potentials. *Neuropsychologia* 119, 330–339
- Juan, C.-H. et al. (2008) Segregation of visual selection and saccades in human frontal eye fields. Cereb. Cortex 18, 2410–2415
- 116. Huddleston, W.E. et al. (2021) Distinct saccade planning and endogenous visuospatial attention maps in parietal cortex: a basis for functional differences in sensory and motor attention. *Cortex* 137, 299–304
- Moore, T. and Zirnsak, M. (2017) Neural mechanisms of selective visual attention. Annu. Rev. Psychol. 68, 47–72
- Super, H. et al. (2004) Correspondence of presaccadic activity in the monkey primary visual cortex with saccadic eye movements. Proc. Natl. Acad. Sci. 101, 3230–3235
- 119. Moore, T. et al. (1998) Visual representations during saccadic eye movements. Proc. Natl. Acad. Sci. U. S. A. 95, 8981–8984
- Tolias, A.S. *et al.* (2001) Eye movements modulate visual receptive fields of V4 neurons. *Neuron* 29, 757–767

- Saber, G.T. et al. (2015) Saccade planning evokes topographically specific activity in the dorsal and ventral streams. *J. Neurosci.* 35, 245–252
- Moore, T. and Chang, M.H. (2009) Presaccadic discrimination of receptive field stimuli by area V4 neurons. *Vis. Res.* 49, 1227–1232
- Steinmetz, N.A. and Moore, T. (2014) Eye movement preparation modulates neuronal responses in area V4 when dissociated from attentional demands. *Neuron* 83, 496–506
- Mitchell, J.F. et al. (2009) Spatial attention decorrelates intrinsic activity fluctuations in macaque area V4. Neuron 63, 879–888
- Cohen, M.R. and Maunsell, J.H.R. (2009) Attention improves performance primarily by reducing interneuronal correlations. *Nat. Neurosci.* 12, 1594–1600
- 126. Rabinowitz, N.C. *et al.* (2015) Attention stabilizes the shared gain of V4 populations. *Elife* 4, e08998
- Steinmetz, N.A. and Moore, T. (2010) Changes in the response rate and response variability of area V4 neurons during the preparation of saccadic eye movements. *J. Neurophysiol.* 103, 1171–1178
- Ni, A.M. et al. (2018) Learning and attention reveal a general relationship between population activity and behavior. Science 359, 463–465
- McAdams, C.J. and Maunsell, J.H. (1999) Effects of attention on orientation-tuning functions of single neurons in macaque cortical area V4. *J. Neurosci.* 19, 431–441
- David, S.V. et al. (2008) Attention to stimulus features shifts spectral tuning of V4 neurons during natural vision. *Neuron* 59, 509–521
- Connor, C.E. et al. (1996) Responses in area V4 depend on the spatial relationship between stimulus and attention. J. Neurophysiol. 75, 1306–1308
- Womelsdorf, T. *et al.* (2006) Dynamic shifts of visual receptive fields in cortical area MT by spatial attention. *Nat. Neurosci.* 9, 1156–1160
- Klein, B.P. et al. (2014) Attraction of position preference by spatial attention throughout human visual cortex. *Neuron* 84, 227–237
- 134. Hartmann, T.S. et al. (2017) Two types of receptive field dynamics in area V4 at the time of eye movements? Front. Syst. Neurosci. 11, 13
- Zirnsak, M. et al. (2014) Visual space is compressed in prefrontal cortex before eye movements. Nature 507, 504–507
- Mineault, P.J. et al. (2016) Enhanced spatial resolution during locomotion and heightened attention in mouse primary visual cortex. J. Neurosci. 36, 6382–6392
- Aydın, Ç. et al. (2018) Locomotion modulates specific functional cell types in the mouse visual thalamus. Nat. Commun. 9, 4882
- Körding, K.P. *et al.* (2007) Causal inference in multisensory perception. *PLoS One* 2, e943
- 139. Atsma, J. *et al.* (2016) Causal inference for spatial constancy across saccades. *PLoS Comput. Biol.* 12, e1004766
- Nobre, A.C. *et al.* (2004) Orienting attention to locations in perceptual versus mental representations. *J. Cogn. Neurosci.* 16, 363–373
- Sergent, C. et al. (2011) Top-down modulation of human early visual cortex after stimulus offset supports successful postcued report. J. Cogn. Neurosci. 23, 1921–1934
- Pestilli, F. et al. (2011) Attentional enhancement via selection and pooling of early sensory responses in human visual cortex. *Neuron* 72, 832–846
- Dugué, L. et al. (2018) Specific visual subregions of TPJ mediate reorienting of spatial attention. Cereb. Cortex 28, 2375–2390
- Dugué, L. *et al.* (2020) Differential impact of endogenous and exogenous attention on activity in human visual cortex. *Sci. Rep.* 10, 21274
- Fernández, A. and Carrasco, M. (2020) Extinguishing exogenous attention via transcranial magnetic stimulation. *Curr. Biol.* 30, 4078–4084
- Ahrens, M.-M. *et al.* (2019) Both dorsal and ventral attention network nodes are implicated in exogenously driven visuospatial anticipation. *Cortex* 117, 168–181
- 147. Van Ettinger-Veenstra, H.M. *et al.* (2009) fMRI-guided TMS on cortical eye fields: the frontal but not intraparietal eye fields



regulate the coupling between visuospatial attention and eye movements. J. Neurophysiol. 102, 3469–3480

- 148. Parker, S. et al. (2021) Establishing the separable contributions of spatial attention and saccade preparation across tasks with varying acuity demands. J. Exp. Psychol. Hum. Percept. Perform. 47, 172–188
- Yeshurun, Y. and Carrasco, M. (2000) The locus of attentional effects in texture segmentation. *Nat. Neurosci.* 3, 622–627
- 150. Talgar, C.P. *et al.* (2004) Covert attention enhances letter identification without affecting channel tuning. *J. Vis.* 4, 22–31